

Edexcel GCSE (9-1) Maths: need-to-know formulae


www.edexcel.com/gcsemathsformulae

Areas


Rectangle = $l \times w$




Parallelogram = $b \times h$



Triangle = $\frac{1}{2} b \times h$



Trapezium = $\frac{1}{2} (a + b)h$




Circles

Circumference = $\pi \times \text{diameter}$, $C = \pi d$

Circumference = $2 \times \pi \times \text{radius}$, $C = 2\pi r$


Area of a circle = $\pi \times \text{radius squared}$, $A = \pi r^2$



Pythagoras


Pythagoras' Theorem

For a right-angled triangle,
 $a^2 + b^2 = c^2$



Trigonometric ratios (new to F)

$\sin x^\circ = \frac{\text{opp}}{\text{hyp}}$, $\cos x^\circ = \frac{\text{adj}}{\text{hyp}}$, $\tan x^\circ = \frac{\text{opp}}{\text{adj}}$




Quadratic equations

The Quadratic Equation


The solutions of $ax^2 + bx + c = 0$,
where $a \neq 0$, are given by $x = \frac{-b \pm \sqrt{(b^2 - 4ac)}}{2a}$

Volumes

Cuboid = $l \times w \times h$



Prism = area of cross section
 $\times \text{length}$



Cylinder = $\pi r^2 h$



Pyramid =
 $\frac{1}{3} \times \text{area of base} \times h$



Compound measures

Speed

$\text{speed} = \frac{\text{distance}}{\text{time}}$



Density

$\text{density} = \frac{\text{mass}}{\text{volume}}$



Pressure

$\text{pressure} = \frac{\text{force}}{\text{area}}$




Trigonometric formulae

Sine Rule $\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$

Cosine Rule $a^2 = b^2 + c^2 - 2bc \cos A$

Area of triangle = $\frac{1}{2} ab \sin C$



Foundation tier formulae

Higher tier formulae

U679



When you have finished with this folder please recycle it

Pearson is committed to reducing its impact on the environment by using responsibly sourced and recycled paper.

Original origami artwork: Mark Bolitho
Lfd / Naki Kouyoumtzis Ocean Image © 123RF: andreykuzhin